

Prevalence and Predictors of Cognitive Impairment among Aged Males and Females: A Population Based Cross-sectional Study

MANINDER KAUR[†], MANKAMAL KAUR* & DIVJOT KAUR[‡]

¹Department of Anthropology, Panjab University, Chandigarh 160014
E-mail: maninderkaur_1@yahoo.in

KEYWORDS: Cognitive scores. Cognitive performance. Illiteracy. Global cognition.

ABSTRACT: The present cross-sectional study is an attempt to determine the prevalence and predictors of cognitive impairment among 98 males and 100 females of Chandigarh, India. All the participants were between 50 to 70 years of age. A 30-point questionnaire i.e Hindi version of MMSE scale was employed to assess the global cognition. Males with higher cognitive scores were significantly lighter, had less developed circumferential measurements and higher waist/hip ratio than males having lower cognitive scores, whereas an opposite trend was witnessed among females. Multivariate regression analysis predicted that illiteracy and > 3 children among females and vegetarian diet among males were associated with impaired cognitive performance. The identification of such potential risk factors may lead to early diagnosis of individuals vulnerable to severe cognitive deficits in later life.

INTRODUCTION

Aging is an inevitable and irreversible process involving a series of psychological, social and biological changes within an individual (Ruvalcaba, 2018). The impressive increment in life expectancy due to the advent of medical facilities has elevated the incidences of various age associated diseases like cardiovascular diseases, hypertension, diabetes and cognitive impairment. Normal cognitive functioning involves proper performance of various activities of human brain like memory, visuo-spatial skills, language, learning and ability of decision making as well as judgment (Sharma *et al.*, 2013). Whereas cognitive impairment is a psychological condition when an individual has trouble in memorizing names or any recently learnt information, face difficulty in learning, concentration and in decision making, and such changes adversely interfere with the routine activities of life. These age-related cognitive alterations exhibit variability in different cognitive

domains as well as in individuals (Glisky, 2007) with the intensity of cognitive decline ranging from mild to severe cases. Review of previous literature (Werner and Korczyn, 2008; Bennett and Thomas, 2014; Pan *et al.*, 2015) have also delineated that cognitive impairment is accompanied by adverse psychiatric conditions like depression, anxiety and sleeplessness, resulting in poor mental as well as physical health, therefore affecting the quality of life of aged people.

The literature has documented many factors such as socio-economic factors (Wu *et al.*, 2016), health conditions (Pektzer and Mafuya, 2012), level of education (Maharani and Tampubolon, 2016), rural or urban setting (Miu *et al.*, 2016), quality of life (Konagaya *et al.*, 2009), low level of depression (Hirsch *et al.*, 2011), high blood pressure (Gao *et al.*, 2009), physical activity (Vance *et al.*, 2005) and healthy diet (Ferdous *et al.*, 2010) that may contribute to the subsequent development of cognitive impairment. Findings of Cronk *et al.* (2010) noticed an inverse association between BMI and cognitive performance. Whereas contrasting view was depicted by Kim *et al.* (2016) and Smith *et al.* (2014) stating a protective

[†] Assistant Professor

* Research scholar, corresponding author

[‡] Student

nature of obesity towards cognitive performance. Studies conducted by Xu *et al.*, (2011) and Whitmer *et al.* (2005) ascertained a positive association between midlife obesity and cognitive decline in later stage of life. Therefore, no uniform conclusion exists regarding the association of increased adiposity and cognitive performance among researchers globally, hence further investigations are required to explore this association.

The reason of association between obesity and cognitive performance is very complex. Various earlier studies on brain imaging have provided the evidence that obese individuals exhibit some morphological changes in their brain such as lower brain volume (Ward *et al.*, 2005) and decreased level of grey matter (Shefer *et al.*, 2013) and white matter (Verstynen *et al.*, 2012), which are responsible for accelerated aging of the brain. The obese individuals exhibit a diminished secretion of adiponectin by adipose tissues (Arnoldussen *et al.*, 2014) thereby reducing the neural protection and energy metabolism (Kaser *et al.*, 2007). These functional and morphological alterations in brain due to adiposity are considered to be the important factor for alterations in cognitive functioning of an individual.

A cross-sectional study conducted by Mohan *et al.* (2019) estimated that the occurrence of mild cognitive impairment amongst older adults of Kerala as 26.06%. Similarly, Singh *et al.*, 2010, documented the overall prevalence of cognitive impairment among population of Bikaner as 19.26% and they further highlighted that this prevalence was higher among females. The conversion rate of mild cognitive impairment to severe condition i.e Alzheimer disease was noted to be approximately 13% (Mridula *et al.*, 2008). Findings of Livingston *et al.* (2017) reported that there can be 35% decrement in occurrence of dementia (severe stage of mild cognitive impairment), if its risk factors can be identified and controlled at its initial stage. Therefore, early recognition of the risk factors that enhances the chances of cognitive impairment among older individuals is important. So, the present study provides us an opportunity to understand the prevalence as well as factors associated with cognitive impairment among adult males and females.

RESEARCH METHODOLOGY

The present cross-sectional study was conducted on 198 participants including 100 females and 98 males. The aim and purpose of the study was explained to all the participants and their verbal consent was taken before conducting the study. The study included healthy individuals aged between 50 to 70 years. Whereas participants with the history of any type of chronic disease, prolonged illness, or any physical deformity were excluded from the present study.

An interview-based schedule was used to gather information like age, marital status, occupation, level of education, dietary habit (vegetarian or non-vegetarian diet), milk intake (number of glasses per day), number of children (≤ 3 or more than 3) and physical activity i.e No activity, walking (3-5 days per week) and regular exercise (5-7 days of yoga, exercise or gym per week).

Anthropometric measurements like height (cm), weight (kg), waist-circumference(cm) and hip circumference(cm) of all the participants were taken using the standard techniques of Weiner and Lourie (1969) and were used to assess their body mass index and waist-hip ratio respectively. All the participants were classified into four categories on the basis of their BMI i.e. underweight (BMI < 18.5 kg/m²), normal weight (18.5-24.9 kg/m²), over weight (25-29.9 kg/m²) and obese (BMI ≥ 30 kg/m²) (WHO, 2008). The presence of central obesity was regarded beyond the cut-off point of waist-hip ratio ≥ 0.90 and ≥ 0.85 for men and women respectively (WHO, 2000).

A 30-point questionnaire i.e. Hindi version of Mini- Mental State Examination scale (Ganguli *et al.*, 1995) was employed to assess the global cognition of all the participants. Individuals scoring below or equal to 23 points were considered as cognitively impaired and the subjects who scored more than 23 points were designated as normal (Tsolaki *et al.*, 2000).

The analysis of collected data was carried out using Statistical Package for the Social Sciences (SPSS) version 16. The quantitative and qualitative data were expressed in mean \pm SD and in percentages (%) or frequencies respectively. Student's t-test was used to compare the mean of continuous variables between individuals of normal cognition and

cognitive impairment groups. Univariate as well as multivariate regression analysis was done to estimate the odds ratio with 95% of confidence Interval (CI) for the various predictors of cognitive impairment in both the sexes. For all the above analysis the level of significance (p value) was considered to be <0.05.

RESULTS

Table 1 presents the general characteristics of 98 males and 100 females of Chandigarh. The mean age

of males and females of present study was 59.34 ± 5.94 years and 58.59 ± 5.83 years respectively. Out of the total sample 73.5% males were working, whereas 73% females were homemakers. The educational status of the participants included in the study demonstrated that most of the males (43.9%) and females (43%) were graduate or more and studied upto metric respectively. On the basis of their cognitive scores participants were categorised into two groups i.e. 41.4% had normal cognition, whereas 58.6% presented impaired cognition.

TABLE 1
General characteristics of the male and female participants

Category		Total (N=198)	Males (N=98)	Females (N=100)
Occupation	Age in years (Mean± SD)	58.96±5.88	59.34±5.94	58.59±5.83
	Home makers N (%)	77 (38.9%)	4(4.1%)	73 (73%)
	Working N (%)	98 (49.5%)	72(73.5%)	26 (26%)
	Retired N (%)	23 (11.6%)	22(22.4%)	1 (1%)
Education	Illiterate N (%)	39 (19.7%)	16 (16.3%)	23(23%)
	Upto matric N (%)	82 (41.4%)	39 (39.8%)	43 (43%)
	Graduation and more	77 (38.9%)	43 (43.9%)	34 (34%)
Marital Status	Married N (%)	192 (97%)	98 (100%)	94 (94%)
	Widow/widower N (%)	6 (3.0%)	0 (0%)	6 (6%)
No. of children	≤3	153 (77.3%)	73(74.5%)	80(80%)
	>3	45 (22.7%)	25(25.5%)	20(20%)
Diet	Vegetarian N (%)	111 (56.1%)	50 (51%)	61 (61%)
	Non-vegetarian N (%)	87 (43.9%)	48 (49%)	39 (39%)
Milk Intake	No	23 (11.6%)	15 (15.3%)	8 (8%)
	1 glass/ day	109 (55.1%)	49 (50%)	60 (60%)
	2 glasses/ day	58 (29.3%)	32 (32.7%)	26 (26%)
	More than 2 glasses/ day	8 (4.0%)	2 (2.0%)	6 (6%)
Physical activity	No activity N (%)	14 (7.1%)	5(5.1%)	9 (9%)
	Walking N (%)	136 (68.7%)	63 (64.3%)	73 (73%)
	Regular exercise N (%)	48 (24.2%)	30 (30.6%)	18 (18%)
Body mass index (BMI)	Under weight (≤18.5 kg/m ²)	04 (2.0%)	2 (2%)	2 (2%)
	Normal (18.5–24.9 kg/m ²)	39 (19.7%)	20 (20.4%)	19 (19%)
	Overweight (≥25.0 kg/m ²)	89 (44.9%)	48 (49%)	41 (41%)
	Obese (≥ 30.0 kg/m ²)	66 (33.3%)	28 (28.6%)	38 (38%)
Cognitive functioning	Normal Cognition N (%)	82 (41.4%)	34 (34.7%)	48 (48%)
	Cognitive impairment N (%)	116 (58.6%)	64 (65.3%)	52 (52%)

The descriptive statistics of various anthropometric variables of males and females with respect to their normal as well as impaired cognitive scores is depicted in Table 2. In the present study cognitive impairment was identified at the prevalence of 65.3% in males and 52% in females. Both the males as well as females with greater cognitive score (MMSE ≥ 24) were taller than their counterparts with lower cognitive scores (MMSE ≤ 23), although differences

were statistically significant only in males. Males with higher cognitive scores were significantly lighter, had less developed circumferential measurements and higher waist/hip ratio than the participants with lower cognitive scores, interestingly an inverse trend was recorded among females for these measurements. The statistically significant differences were observed for waist circumference in both the sexes and for BMI in females only.

TABLE 2

Descriptive statistics of anthropometric variables of males and females with respect to their cognitive scores

Anthropometric and lifestyle Characteristics	Males (N=98)		t-test	Females (N=100)		t-test
	Normal cognitive scores	Cognitive impairment		Normal cognitive scores	Cognitive impairment	
	MMSE \geq 24 (n=34)	MMSE \leq 23 (n=64)		MMSE \geq 24 (n=48)	MMSE \leq 23 (n=52)	
Height (cm)	164.74 \pm 4.97	163.82 \pm 52.31	0.000***	159.03 \pm 5.25	158.32 \pm 6.73	1.995
Weight (kg)	72.88 \pm 10.10	73.13 \pm 10.00	0.006*	70.38 \pm 12.42	69.56 \pm 11.17	0.973
BMI	28.02 \pm 3.99	28.05 \pm 4.56	0.762	28.74 \pm 5.37	28.35 \pm 5.97	0.035*
Waist circumference	93.76 \pm 15.04	96.02 \pm 12.45	0.000***	93.53 \pm 13.08	90.38 \pm 15.71	0.009*
Hip circumference	94.01 \pm 12.77	98.15 \pm 12.66	0.396	102.09 \pm 14.44	99.06 \pm 12.54	0.693
Waist/hip ratio	0.99 \pm 0.07	0.97 \pm 0.05	0.931	0.91 \pm 0.05	0.92 \pm 0.04	1.387

Level of significance = $p \leq 0.05^*$, $p \leq 0.01^{**}$, $p \leq 0.001^{***}$

Figure 1 outlines the mean MMSE scores with respect to different categories of body mass index in both males and females of the present study. It is inferred from the figure that obese males (MMSE scores= 22.39) had highest mean global MMSE score followed by overweight (MMSE scores= 22.08) and normal weight (MMSE scores= 21.85) individuals and

lowest scores were observed among underweight males (MMSE scores = 19.00). Whereas among females the mean cognitive score was highest (MMSE scores= 22.17) among overweight females, followed by obese (MMSE scores= 22), underweight (MMSE scores = 21) and normal weight (MMSE scores= 20.84) subjects.




Figure 1 Mean MMSE scores with respect to different categories of body mass index in both males and females

Univariate regression analysis was conducted to estimate the possible predictors of cognitive impairment among males and females. The results of the univariate regression analysis (Table 3) depicted that out of the various factors only level of education and number of children were the significant predictors of cognitive impairment among females. Women having >3 children exhibited higher chances of cognitive decline with odds ratio 27.06 (95% CI 3.45 – 212.22, $p \leq 0.05$). Similarly, illiteracy was also significantly associated with greater chances of cognitive impairment with odds ratio of 34.46 (95% CI

4.41-269.26, $p \leq 0.01$). Whereas among males vegetarian diet exhibited 0.34 folds (95% CI 0.14-0.83, $p \leq 0.05$) higher chances of cognitive impairment. The variables exhibiting statistically significant associations in the univariate model were further subjected to multivariate regression model.

According to multivariate regression model (Table 4), illiterate women were found to have a 27.5 folds (95% CI 3.38-223.44, $p < 0.001$) higher probability of the cognitive impairment. Women having >3 children were 20.64 times (2.48-171.66, $p < 0.001$) more likely to be associated with impaired cognitive

performance. Vegetarian males were found to present 0.34 folds (95% CI 0.14-0.83, $p < 0.01$) higher chances of the cognitive impairment.

TABLE 3

Univariate analysis of factors associated with cognitive impairment among males and females

Variables	Males (N=98)		Females (N=100)	
	Odds ratio (95% CI)	p value	Odds ratio (95% CI)	p value
Age (In years) ^a	≤ 55	1	1	
	56-64	0.609(0.230-1.609)	1.235 (0.507-3.010)	0.6420.083
	≥65	0.727(0.219-2.411)	2.647 (0.880-7.966)	
Level of Education	Illiterate	4.48(0.95-21.03)	34.46(4.41-269.26)	0.001**
	Literate	1	1	
No. of Children	≤3	1	1	
	>3	1.50(0.55-4.07)	27.06(3.45-212.22)	0.002*
Diet	Vegetarian	0.34 (0.14-0.83)	0.78(0.34-1.76)	0.556
	Non-vegetarian	1	1	
Physical Activity	No activity	0.43(0.04-4.36)	4.37(0.70-27.16)	0.113
	Waking	0.92(0.37-2.29)	1.28(0.45-3.62)	0.636
	Regular exercise/ yoga	1	1	
Hypertension	Yes	0.66 (0.21-2.10)	1.57(0.55-4.45)	0.395
	No	1	1	
Body Mass Index	Underweight	0.42(0.02-8.04)	0.72(0.03-13.45)	0.831
	Normal weight	1	1	
	Overweight	0.65(0.21-2.00)	0.69(0.23-2.07)	0.512
	Obese	1.07(0.30-3.77)	0.80(0.26-2.45)	0.707
Waist-Hip ratio	Normal	1	1	0.110
Abdominal obesity (WHR ≥0.90 for men and WHR >0.85 for women)		0.69(0.29-1.64)	5.93(0.66-52.72)	

Level of significance = $p \leq 0.05^*$, $p \leq 0.01^{**}$, $p \leq 0.001^{***}$

TABLE 4

Multivariate analysis of factors associated with cognitive impairment among males and females

Variables	Odds Ratio (95% CI)	p-value
<i>Females: Model 1</i>		
Level of education		
illiterate	27.5(3.38-223.44)	0.00**
Literate	1	
Number of Children		
>3	20.64(2.48-171.66)	0.00**
≤3	1	
<i>Males: Model 1</i>		
Diet		
Vegetarian	0.34 (0.14-0.83)	0.01*
Non-vegetarian	1	

Level of significance = $p \leq 0.05^*$, $p \leq 0.01^{**}$, $p \leq 0.001^{***}$

DISCUSSION

In the present sample overall prevalence of cognitive impairment was estimated as high as 58.6%, with higher prevalence among males (65.3%) as compared to their female (52%) counterparts. Similarly, Ramchandran *et al.* (2018) also conducted a cross-sectional study to analyse the cognitive functioning using MMSE Scale on 100 elderly subjects of Kerala and observed the prevalence of cognitive impairment

to be 55%. Previous cross-sectional studies recorded 42.9% cognitive impairment in Varanasi (Gambhir *et al.*, 2014), 38% in Hyderabad (Rao *et al.*, 2014) and 42.7% in Chennai (Samuel *et al.*, 2016). Recently, Vidyanti *et al.* (2020) documented the higher prevalence of cognitive impairment (75.5%) among people of Indonesia as assessed by Indonesian version of Montreal Cognitive Assessment Scale. Similar results were documented by Arjuna *et al.* (2017)

who indicated its prevalence to be about 59% and 80% among urban and rural population of Indonesia respectively. Earlier authors (Wilkie and Eisdorfer, 1971) considered that cognitive impairment is not a product of normal aging process, but it should be viewed as secondary to some pathologic conditions.

In the present study prevalence of cognitive impairment was higher among males than females. Recently Abi-Ghanem *et al.* (2020) explained the effects of androgens on cerebrovascular function and noticed higher risk of stroke and dementia (VCID) among males as compared to females throughout the lifespan. While conflicting findings were seen in the study of Liu *et al.* (2019) where females exhibited significantly higher prevalence of cognitive impairment than males (males 41.5% vs females 58.5%, $\chi^2 = 9.49^{**}$). In the present study most of the males (73.5%) were working, while substantial percentage of women were involved in household chores only. Therefore, males were having more stressful circumstances than their female counterparts. This could be one of the reasons for higher prevalence of cognitive impairment among males.

The results of the present study depicted that the males with normal cognition were significantly taller, lighter but lower BMI and circumferential measurements than cognitively impaired males. West *et al.* (2015) also observed that men with short stature were associated with poor cognitive functioning especially in the domains of global cognition ($p = 0.016$), executive functioning ($p = 0.001$) and attention ($p = 0.007$). Similar observations were noticed by Quan *et al.* (2013) among Korean population that men with short height exhibited higher chances of cognitive impairment. A report of Perneckzy (2010) explained that individuals with short height have smaller head circumference, resulting in lower brain reserve, which in turn is associated with decline in cognitive functioning. Divergent results were observed by Noh *et al.* (2017) among Korean population, where males with cognitive impairment presented higher values of height, BMI and waist-hip ratio as compared to their cognitively normal counterparts.

In the present study females with normal cognition were taller, heavier, higher BMI and circumferential measurements than cognitively

impaired females. Similar results were observed among three hundred twenty participants of South Korea aged 65 years or older, in which females with normal cognition as assessed by Korean version of Mini Mental state examination exhibited significantly higher body mass index (25.52 ± 3.58) as compared to cognitively impaired females (23.72 ± 3.14) (Noh *et al.* 2017). Konda *et al.* (2018) also conducted a cross-sectional study and reported that both lower weight and waist-hip ratio were significantly associated with higher chances of cognitive impairment among older adults (60 years or older) of Hyderabad, India. Similarly, Skinner *et al.* (2017) evaluated the cognitive performance of three hundred thirty-one individuals (aged 55 years or above), out of which majority (94.6%) of the participants were women and documented the beneficial effect of obesity towards cognitive functioning. Findings of Bagger *et al.* (2004) documented that body fat mass may play a protective role in halting cognitive deterioration and this protection can be attributed to nonprotein-bound estradiol produced by visceral adipocytes.

Health benefits could be accomplished by identifying possible determinants, in the present study univariate regression analysis and further multivariate regression depicted that out of the various factors only illiteracy (OR=27.5, 95% CI 3.38-223.44) and more than three children (OR=20.64, 95% CI 2.48-171.66) were the significant predictors of cognitive impairment among females, whereas vegetarian diet (OR=0.34, 95% CI 0.14-0.83) was significantly associated with lower chances of cognitive impairment among males. In accordance to our findings, Rao *et al.* (2014) also evaluated the cognitive functioning of older adults (aged 60 years or above) of Hyderabad using Mini Mental state Examination scale and observed that covariates such as age, gender (female), lower educational level, marital status (widow/ widower) and nuclear family type were significantly associated with higher chances of cognitive impairment. Similar to these results various other studies (Khairiah *et al.*, 2016; Maroof *et al.*, 2016 and Konda *et al.*, 2018) also observed a positive association between cognitive impairment and no formal education among different elderly populations of Malaysians, rural areas of Aligarh and Urban area of Hyderabad respectively. In Latin America, the illiteracy rate was noted to be 9.3% and the prevalence of

dementia (severe stage of mild cognitive impairment) in illiterate population was almost double as compared to the literates (Nitrini *et al.*, 2009). In India, Sengupta *et al.* (2014) analysed the correlates of impaired cognition among older adults (60 years or above) of Ludhiana, Punjab using Hindi Mental state examination scale and found that older age, no education, widowed status, unemployment and lower monthly income (< 1000 rupees) were the significant predictors of cognitive impairment. A study performed by Grossi *et al.* (1993) illustrated that the basic learning during the formative years of schooling is responsible for improvement of mental strategies, and are well preserved during the normal aging process. Whereas Bruki, (2010) explained that illiteracy is associated with lower cognitive performance, due to unsuitable adaptation of a neuropsychological test for a particular population and difficulty in cognitive evaluation therefore documenting poor cognitive scores among illiterate population.

The major limitation of the current study is its cross-sectional nature, as it does not examine the longitudinal performance of the participants and the true causal relationship could only be identified through extended follow-up studies among both males and females. The study has only assessed the global cognitive functioning of the participants without specifically focusing on each cognitive domain. More information regarding the dietary patterns of the participants would have presented better results.

Despite of certain limitations there are several strengths of the study. To the best of our knowledge this is the first report assessing the prevalence and factors associated with cognitive impairment among both males and females of Chandigarh, India. The study provides us the epidemiological data for a population for which, as yet, there is no such information.

CONCLUSION

With expanding age people are spending more years of their life in old ages. World health organisation (2019) had documented that approximately 10 million new cases suffer from dementia (severe stage of mild cognitive impairment) every year. In addition to this, mild cognitive impairment also leads to psychological symptoms like

depression and insomnia among both patients and caregivers (Na, 2019). In order to flatten this increasing curve, there is an urgent need for the identification of potential risk factors, which can lead to early diagnosis of individuals vulnerable to severe cognitive deficit in later life and is also important for the development of preventive strategies, which can therefore help in halting the progression of cognitive deterioration among elderly population.

REFERENCES CITED

- Abi Ghanem C., L. S. Robison and K. L. Zuloaga 2020. Androgens' effects on cerebrovascular function in health and disease. *Biology of Sex Differences*, 11(1):1-8. Doi: <https://doi.org/10.1186/s13293-020-00309-4>
- Arjuna T., S. Soenen, R. A. Hasnawati, K. Lange, I. Chapman and N. D. Luscombe-Marsh 2017. A cross-sectional study of nutrient intake and health status among older adults in Yogyakarta Indonesia. *Nutrients*, 9(11):1240. Doi: 10.3390/nu9111240
- Arnoldussen I. A., A. J. Kiliaan and D. R. Gustafson 2014. Obesity and dementia: adipokines interact with the brain. *European Neuropsychopharmacology*, 24(12):1982-99. Doi:10.1016/j.euroneuro.2014.03.002
- Bagger Y. Z., L. B. Tanko, P. Alexandersen, G. Qin and C. Christiansen 2004. The implications of body fat mass and fat distribution for cognitive function in elderly women. *Obesity Research*, 12(9):1519-26. Doi: <https://doi.org/10.1038/oby.2004.189>
- Bennett S., and A. J. Thomas 2014. Depression and dementia: cause, consequence or coincidence?. *Maturitas*. 79(2):184-90. Doi: <https://doi.org/10.1016/j.maturitas.2014.05.009>
- Brucki S. M. 2010. Illiteracy and dementia, *Dementia & Neuropsychologia*, 4(3):153-7. [PubMed: 29213680]
- Cronk B. B., D. K. Johnson and J. M. Burns 2010. Alzheimer's Disease Neuroimaging Initiative. Body mass index and cognitive decline in mild cognitive impairment. *Alzheimer disease and associated disorders*, 24(2):126. Doi: 10.1097/WAD.0b013e3181a6bf3f.
- Ferdous T., T. Cederholm, Z. N. Kabir, J. D. Hamadani and A. Wahlin 2010. Nutritional status and cognitive function in community living rural Bangladeshi older adults: data from the poverty and health in ageing project. *Journal of the American Geriatrics Society*, 58(5):919-24. Doi: <http://dx.doi.org/10.1182/03/2394-6040.ijcmph20180788>
- Gambhir I. S., V. Khurana, D. Kishore, A. K. Sinha and S. C. Mohapatra 2014. A clinico-epidemiological study of cognitive function status of community-dwelling elderly. *Indian journal of psychiatry*, 56(4):365. Doi: <https://dx.doi.org/10.4103%2F0019-5545.146531>
- Ganguli M., G. Ratcliff, V. Chandra, S. Sharma, J. Gilby, R.

- Pandav, S. Belle, C. Ryan, C. Baker, E. Seaberg and S. Dekosky 1995. A-Hindi version of the MMSE: the development of a cognitive screening instrument for a largely illiterate rural elderly population in India. *International Journal of Geriatric Psychiatry*, 10(5):367-77. Doi: <https://doi.org/10.1002/gps.930100505>
- Gao S., Y. Jin, F. W. Unverzagt, C. Liang, K. S. Hall, F. Ma, J. R. Murrell, Y. Cheng, J. Matesan, J. Bian and P. Li 2009. Hypertension and cognitive decline in rural elderly Chinese. *Journal of the American Geriatrics Society*, 57(6):1051-7. Doi: <https://doi.org/10.1111/j.1532-5415.2009.02267>.
- Glisky, E. L 2007. *Changes in cognitive function in human aging*. Brain aging. 19:3-20.
- Grossi D., G. Correria, C. Cause, M. A. Ruscitto, V. Vecchione, M. V. Vigliardi and G. Nolfè 1993. Evaluation of the influence of illiteracy on neuropsychological performances by elderly persons. *Perceptual and motor skills*, 77(3):859-66. Doi: <https://doi.org/10.2466/2Fpms.1993.77.3.859>.
- Hirsch J. K., F. M. Sirois and J.M. Lyness 2011. Functional impairment and depressive symptoms in older adults: Mitigating effects of hope. *British Journal of Health Psychology*. 16(4):744-60. Doi: <https://doi.org/10.1111/j.2044-8287.2010.02012.x>
- Kaser S., T. Tatarczyk, K. Salzmann, A. Stadlmayr, B. Iglseider, V. Mackevics, C. Ciardi, J. Engl, C. Röss, M. Laimer and A. Sandhofer 2007. Effect of Obesity and Insulin Sensitivity on Adiponectin Isoform Distribution. *Diabetes*, 2:56. Doi: <https://doi.org/10.1111/j.1365-2362.2008.02028.x>
- Khairiah K., C. S. Mooi and T. A. Hamid 2016. Prevalence and factors associated with mild cognitive impairment on screening in older Malaysians. *Dusunen Adam The Journal of Psychiatry and Neurological Sciences*, 29(4):298. Doi: <https://dx.doi.org/10.5350/DAJPN2016290401>
- Kim S., Y. Kim and S. M. Park 2016. Body mass index and decline of cognitive function. *PLOS One*, 11(2):e0148908.
- Konagaya Y., T. Watanabe, T. Ohta and K. Takata. Relationship between quality of life (QOL) and cognitive function among community-dwelling elderly. *Japanese Journal of Geriatrics*, 46(2):160-7. Doi: 10.3143/geriatrics.46.160
- Konda P. R., P. K. Sharma, A. R. Gandhi and E. Ganguly 2018. Correlates of cognitive impairment among Indian Urban Elders. *Journal Of Gerontology & Geriatric Research*. 7(6). Doi: <https://dx.doi.org/10.4172%2F2167-7182.1000489>
- Liu Z., H. Yang, S. Chen, J. Cai and Z. Huang 2019. The association between body mass index, waist circumference, waist-hip ratio and cognitive disorder in older adults. *Journal of Public Health*, 41(2):305-12. Doi: <https://doi.org/10.1093/pubmed/fdy121>
- Livingston G., A. Sommerlad, V. Orgeta, S. G. Costafreda, J. Huntley, D. Ames, C. Ballard, S. Banerjee, A. Burns, J. Cohen-Mansfield and C. Cooper 2017. Dementia prevention, intervention, and care. *The Lancet*. 390(10113):2673-734. Doi: [https://doi.org/10.1016/S0140-6736\(17\)31363-6](https://doi.org/10.1016/S0140-6736(17)31363-6)
- Maharani A., and G. Tampubolon 2016. National economic development status may affect the association between central adiposity and cognition in older adults. *PloS One*, 11(2):e0148406. <https://doi.org/10.1371/journal.pone.0148406>
- Maroof M., A. Ahmad, N. Khalique, M. A. Ansari, M. S. Shah and U. Eram 2016. Prevalence and determinants of cognitive impairment among rural elderly population of Aligarh. *National Journal of Community Medicine*, 7(3):189-92.
- Miu J., J. Negin, A. Salinas-Rodriguez, B. Manrique-Espinoza, A. L. Sosa-Ortiz, R. Cumming and P. Kowal 2016. Factors associated with cognitive function in older adults in Mexico. *Global Health Action*. 9(1):30747. Doi: <https://doi.org/10.3402/gha.v9.30747>
- Mohan D., T. Iype, S. Varghese, A. Usha and M. Mohan 2019. A cross-sectional study to assess prevalence and factors associated with mild cognitive impairment among older adults in an urban area of Kerala, South India. *British Medical Journal*, 9(3):e025473. Doi: <http://dx.doi.org/10.1136/bmjopen-2018-025473>
- Mridula K. R., A. Suvarna, M. Shailaja, C. S. Kumari and S. Kaul 2008. Mild Cognitive Impairment: Characterisation of clinical, neuropsychological, imaging profile and outcome. *Annals of Indian Academy of Neurology*. 11.
- Na K. S. 2019. Prediction of future cognitive impairment among the community elderly: a machine-learning based approach. *Scientific Reports*. 9(1):1-9. Doi: <https://dx.doi.org/10.1038%2Fs41598-019-39478-7>
- Nitrini R., C. M. Bottino, C. Albala, N. S. Capuñay, C. Ketzoian, J. J. Rodriguez, G. E. Maestre, A. T. Ramos-Cerqueira and P. Caramelli 2009. Prevalence of dementia in Latin America: a collaborative study of population-based cohorts. *International Psychogeriatrics*, 21(4):622-30. Doi: <http://dx.doi.org/10.1017/S1041610209009430>
- Noh H. M., S. Oh, H. J. Song, E. Y. Lee, J. Y. Jeong, O. H. Ryu, K. S. Hong and D. H. Kim 2017. Relationships between cognitive function and body composition among community-dwelling older adults: a cross-sectional study. *BMC Geriatrics*, 17(1):1-9. Doi: 10.1186/s12877-017-0651-9
- Pan C.W., X. Wang, Q. Ma, H. P. Sun, Y. Xu and P. Wang 2015. Cognitive dysfunction and health-related quality of life among older Chinese. *Scientific Reports*, 5(1):1-8. Doi: 10.1038/srep17301
- Peltzer K., and N. Phaswana-Mafuya 2012. Cognitive functioning and associated factors in older adults in South Africa. *South African Journal of Psychiatry*, 2012 18(4):157-63. <https://doi.org/10.7196/sajp.368>
- Perneczky R., S. Wagenpfeil, K. L. Lunetta, L. A. Cupples,

- R. C. Green, C. DeCarli, L. A. Farrer and A. Kurz 2010. Head circumference, atrophy, and cognition: implications for brain reserve in Alzheimer disease. *Neurology*, 75(2):137-42. Doi: <http://doi.org/10.1212/WNL.0b013e3181e7ca97>
- Quan S. A., J. Y. Jeong and D. H. Kim 2013. The relationship between height and cognitive function among community-dwelling elderly: Hallym Aging Study. *Epidemiology and Health*. 35.
- Ramachandran R., J. M. Mundodan, C. R. Saju and V. M. Joshy 2018. Nutritional status and cognitive impairment in elderly population in a rural area of Thrissur district, Kerala. *International Journal of Community Medicine and Public Health*, 5:1218-23. Doi: <http://dx.doi.org/10.18203/2394-6040.ijcmph20180788>
- Ramachandran V., and P. Aryani 2018. Association between educational level and hypertension with decrease of cognitive function among elderly at Puskesmas Mengwi I. *Intisari Sains Medis*, 9(1):43-8. Doi: 10.15562/ism.v9i1.154
- Rao S. S., S. K. Chennamsetty and S. R. Kuna 2014. A cross-sectional study of cognitive impairment and morbidity profile of inmates of old age home. *Scholars Journal of Applied Medical Sciences*, 2(4):1506-3. Corpus ID:20242479
- Ruvalcaba M., M. Neyda, E. D. Arias-Merino, M.E. Flores-Villavicencio, M. Rodríguez-Díaz, I. F. Díaz-García 2018. Cognitive Aging. *Gerontology*, 4:143. Doi:10.5772/intechopen.71551.
- Samuel R., C. S. McLachlan, U. Mahadevan and V. Isaac 2016. Cognitive impairment and reduced quality of life among old-age groups in Southern Urban India: home-based community residents, free and paid old-age home residents. *QJM: An International Journal of Medicine*, 109(10):653-9. Doi: <https://doi.org/10.1093/qjmed/hcw040>
- Sengupta P., A. I. Benjamin, Y. Singh and A. Grover 2014. Prevalence and correlates of cognitive impairment in a north Indian elderly population. *WHO South-East Asia Journal of Public Health*, 3(2):135-43. Doi: 10.4103/2224-3151.206729.
- Sharma D., S. R. Mazta, and A. Parashar 2013. Prevalence of cognitive impairment and related factors among elderly: A population-based study. *Journal of Dr. NTR University of Health Sciences*, 2(3):171. Doi: 10.4103/2277-8632.117182.
- Shefer G., Y. Marcus and N. Stern 2013. Is obesity a brain disease?. *Neuroscience & Biobehavioral Reviews*. 37(10):2489-503. Doi: <https://doi.org/10.1016/j.neubiorev.2013.07.015>
- Singh V. B 2010. Prevalence of mild cognitive impairment in elderly population of North India.
- Skinner J. S., W. M. Abel, K. McCoy and C. H. Wilkins 2017. Exploring the "obesity paradox" as a correlate of cognitive and physical function in community-dwelling black and white older adults. *Ethnicity & Disease*. 27(4):387. Doi: <https://dx.doi.org/10.18865%2Fed.27.4.387>
- Smith E., P. E. Bailey, J. Crawford, K. Samaras, B. T. Baune, L. Campbell, N. Kochan, J. Menant, D. L. Sturnieks, H. Brodaty and P. Sachdev P. Adiposity Estimated Using Dual Energy X Ray Absorptiometry and Body Mass Index and Its Association with Cognition in Elderly Adults. *Journal of the American Geriatrics Society*, 62(12):2311-18. Doi: <https://doi.org/10.1111/jgs.13157>
- Tsolaki M., V. Iakovidou, H. Navrozidou, M. Aminta, T. Pantazi and Kazis A 2000. Hindi Mental State Examination (HMSE) as a screening test for illiterate demented patients. *International journal of geriatric psychiatry*, 15(7):662-4. Doi: [https://doi.org/10.1002/1099-1166\(200007\)15:7%3C662::AID-GPS171%3E3.0.CO;2-5](https://doi.org/10.1002/1099-1166(200007)15:7%3C662::AID-GPS171%3E3.0.CO;2-5)
- Vance D. E., V. G. Wadley, K. K. Ball, D. L. Roenker and M. Rizzo 2005. The effects of physical activity and sedentary behaviour on cognitive health in older adults. *Journal of Aging and Physical Activity*, 13(3):294-313. Doi: <https://doi.org/10.1123/japa.13.3.294>
- Verstynen T. D., A. M. Weinstein, W. W. Schneider, J. M. Jakicic, D. L. Rofey and K. I. Erickson 2012. Increased body mass index is associated with a global and distributed decrease in white matter microstructural integrity. *Psychosomatic Medicine*, 74(7):682. Doi: <https://dx.doi.org/10.1097%2FPSY.0b013e318261909c>
- Vidyanti A. N., M. Hardhantyo, B. S. Wiratama, A. Prodjohardjono and C. J. Hu 2020. Obesity is less frequently associated with cognitive impairment in elderly individuals: a cross-sectional study in Yogyakarta, Indonesia. *Nutrients*, 12(2):367. Doi: <https://doi.org/10.3390/nu12020367>
- Werner, P., and A.D. Korczyn 2008. Mild cognitive impairment: conceptual, assessment, ethical, and social issues. *Clinical Interventions In Aging*, 3(3):413. Doi: <https://dx.doi.org/10.2147%2Fcia.s1825>
- Ward M. A., C. M. Carlsson, M. A. Trivedi, M. A. Sager and S. C. Johnson 2005. The effect of body mass index on global brain volume in middle-aged adults: a cross sectional study. *BMC Neurology*, 5(1):1-7. Doi:10.1186/1471-2377-5-23.
- Weiner, J. S. and J. A. Lourie 1969. *Human biology, a guide to field methods*. International Biological Program Handbook No. 9 Blackwell, Oxford.
- West R. K., R. Ravona-Springer, A. Heymann, J. Schmeidler, D. Leroith, K. Koifman, E. Guerrero-Berroa, R. Preiss, H. Hoffman, J. M. Silverman and M. S. Beeri 2015. Shorter adult height is associated with poorer cognitive performance in elderly men with type II diabetes. *Journal of Alzheimer's Disease*. 44(3):927-35.
- Whitmer R. A., E. P. Gunderson, E. Barrett-Connor, C. P. Quesenberry and K. Yaffe 2005. Obesity in middle age and future risk of dementia: a 27 year longitudinal population based study. *British Medical Journal*, 330(7504):1360. Doi: 10.1136/bmj.38446.466238.E0.

- Wilkie F., and C. Eisdorfer 1971. Intelligence and blood pressure in the aged. *Science*, 172(3986):959-62. Doi: 10.1126/science.172.3986.959
- World Health Organization 2000. The Problem of Obesity. [Available from: [http://www.who.int/trs/WHO_TRS_894_\(part1\).pdf](http://www.who.int/trs/WHO_TRS_894_(part1).pdf)].
- World Health Organization 2008. Waist Circumference and Waist-Hip Ratio. Report of WHO Expert Consultation.
- World Health Organization 2019. <http://www.who.int/news-room/fact-sheets/detail/dementia>
- Wu F., Y. Guo, Y. Zheng, W. Ma, P. Kowal, S. Chatterji, L. Wang 2016. Social-economic status and cognitive performance among Chinese aged 50 years and older. *Plos One*, 11(11):e0166986. Doi:10.1371/journal.pone.0166986
- Xu W. L., A. R. Atti, M. Gatz, N. L. Pedersen, B. Johansson and L. Fratiglioni 2011. Midlife overweight and obesity increase late-life dementia risk: a population-based twin study. *Neurology*, 76(18):1568-74. Doi: 10.1212/WNL.0b013e3182190d09